arXiv:0807.3224v1 [cond-mat.other] 21 Jul 2008

Observation of long-lived polariton states in semiconductor microcavities across the parametric threshold

D. Ballarini,[1] D. Sanvitto,[1] A. Amo,[1] L. Viña,[1] M. Wouters,[2] I. Carusotto,[3] A. Lemaitre,[4] and J. Bloch[4]

[1] *Dep. Física de Materiales, Univ. Autonóma de Madrid, 28049 Madrid, Spain*
[2] *Institute of Theoretical Physics, Ecole Polytechnique Fédérale de Lausanne EPFL, CH-1015 Lausanne, Switzerland*
[3] *BEC-CNR-INFM and Dipartimento di Fisica, Università di Trento, I-38050 Povo, Italy*
[4] *LPN/CNRS, Route de Nozay, 91460, Marcoussis, France*

The excitation spectrum around the pump-only stationary state of a polariton optical parametric oscillator (OPO) in semiconductor microcavities is investigated by time-resolved photoluminescence. The response to a weak pulsed perturbation in the vicinity of the idler mode is directly related to the lifetime of the elementary excitations. A dramatic increase of the lifetime is observed for a pump intensity approaching and exceeding the OPO threshold. The observations can be explained in terms of a critical slowing down of the dynamics upon approaching the threshold and the following onset of the soft Goldstone mode.

PACS numbers:

One of the most striking consequences of the quantum nature of matter is Bose-Einstein condensation, a phase transition that does not depend on interactions between the microscopic constituents but is driven by their indistinguishability. A quite general property of Bose-Einstein condensed quantum fluids is superfluidity, i.e. the ability to flow through a container almost without any friction. Since the first discovery in 4-Helium, this effect was investigated in a variety of quantum fluids, ranging from 3-Helium to ultracold atomic gases **(author?)** [1]. Inspired by recent developments in the study of phase transitions in non equilibrium systems **(author?)** [2], researchers are presently working on the extension of the quantum fluid concept to many particle systems whose state is no longer determined by a thermodynamical equilibrium condition, but rather by a balance between the external driving and dissipation.

In this perspective, exciton-polaritons in semiconductor microstructures appear as very promising objects, as their mixed excitonic and photonic nature allows for a number of remarkable properties, e.g. a very light mass, significant interactions, and the possibility of all-optical manipulation and diagnostics **(author?)** [3]. On the other hand, the finite lifetime of polaritons requires some pumping mechanism to continuously replenish the system, which forbids the establishment of a true thermal equilibrium in the gas. Several groups have reported the observation of spontaneous coherence in spatially extended systems of polaritons far from thermodynamical equilibrium with mechanisms that can be interpreted as non-equilibrium analogs of Bose-Einstein condensation (BEC) **(author?)** [4].

In spite of the different pumping schemes (resonant for the OPO case **(author?)** [5, 6, 7, 8], non-resonant for the so-called polariton BEC case **(author?)** [9]), a $U(1)$ symmetry is spontaneously broken in all cases and coherence is not simply inherited from the pump beam. Away from the bistability regime **(author?)** [10], the OPO transition is smooth and shows a critical behavior that closely resembles the one of a second-order like phase transition **(author?)** [11]. A typical feature of second order phase transitions is the critical slowing down of the decay rate of the elementary excitations upon approaching the transition from below. When a continuous symmetry is spontaneously broken above the transition point, the lifetime of elementary excitations remains divergent in the long wavelength limit in agreement with the Goldstone theorem of statistical mechanics **(author?)** [12, 13].

In the present Letter, we experimentally investigate and theoretically model the lifetime of the elementary excitations of a quantum fluid of polaritons as the pump intensity is spanned across the threshold in the optical parametric oscillator (OPO) configuration. The steady state of the system is probed by injecting extra polaritons by means of a weak pulsed beam, and the decay time of the response is measured as a function of the pump intensity. A dramatic slowing down of the dynamics is observed as the threshold is approached from below: close to the threshold, the decay time can become orders of magnitude longer than the typical life time of polaritons, and it remains very long even well above the threshold. Good agreement between the experimental observations and the theoretical model based on the generalized polariton Gross-Pitaevskii equation is found, which supports the present theoretical understanding of the strongly modified dispersion of the elementary excitations in presence of a strong pump beam. In particular, this observation suggests the possibility of investigating the polariton dynamics beyond the limits imposed by the intrinsic polariton life time.

The microcavity sample used in these experiments is a GaAs/AlAs-based $\lambda/2$ cavity with a top (bottom) Bragg mirror of 15 (25) $Al_{0.1}Ga_{0.9}As$/AlAs pairs, grown on a GaAs substrate. A 20 nm wide GaAs quantum well is embedded at the antinode position of the cavity mode. The sample is kept at a constant temperature of 10 K.

The quantum well excitons are in strong coupling with the cavity mode, with a Rabi splitting of $2\Omega_R = 4.4$ meV. All the experiments reported here are performed in the resonance region, where the $k = 0$ cavity-mode has the same energy as the exciton state.

In the OPO configuration, polaritons are coherently injected into the microcavity by a pump beam, which resonantly populates a polariton mode with a defined momentum and energy (pump state). Our pump beam is a continuous-wave laser source (Ti:Al_2O_3), which excites the sample with an incident angle of 10° and has a 45 μm spot diameter. Its frequency is chosen close to resonance with the lower polariton branch (LPB) in a way to inject polaritons with a given wave vector k_p and energy $\hbar\omega_p$. Polariton-polariton binary collisions are responsible for the parametric scattering of pump polaritons into a pair of distinct signal and idler modes. The efficiency of the parametric processes was optimized by tuning the pump at a frequency $\hbar\omega_P = 1.5273$ eV, slightly above the linear-regime LP dispersion **(author?)** [14]**(author?)** [10], $\varepsilon_{LPB}(k_P)$, $\hbar\omega_p - \varepsilon_{LPB}(k_P) \sim \gamma(k_p)$. Here $\gamma(k_p) \sim 0.4$ meV is the LPB linewidth at k_p, while the laser linewidth is around 0.1 meV. An additional 2 ps-long *probe* pulse, coming from a different laser source with a repetition rate of 82 MHz, is incident on the sample with a tunable angle and is focused within the pump spot with a smaller, 25 μm, spot diameter. Photoluminescence (PL) is collected and analyzed by a spectrograph coupled either to a streak- or a conventional CCD-camera. The emission in the far field is visualized by means of a lens on the Fourier plane $(k_x; k_y)$: a direction k_y of this plane is selected and energy-resolved by the spectrograph, which allows the direct observation on the CCD camera of any two-dimensional section $(k_x; E)$ of the polariton dispersion. To obtain time-resolved PL images a streak camera is used in either a $(E; t)$ or a $(k_x; t)$ configuration, which allows for a complete study of the dynamics at different energies and momenta.

Firstly, we have investigated the stationary-state polariton emission in the absence of the probe. Typical energy-momentum emission patterns are shown in Fig.1 for two different values of the CW pump power I_p. Although the pump wavevector lies well outside the k-space region imaged in Fig.1, a small polariton occupation of the LPB bottom still appears as a consequence of incoherent relaxation processes even at low pump powers (lower panel). As the polariton density is very low, the photoluminescence spectrum is concentrated on the linear-regime LPB branch.

At higher pump intensities, polariton-polariton interactions are able to significantly modify the emission pattern and, in particular, are responsible for parametric processes, where two pump polaritons at k_p are transformed into a pair of signal and idler polaritons of wavevectors $k_{s,i}$, respectively. Thanks to the large size of the pump spot, the wavevector is approximately conserved, $k_s \simeq 2\,k_p - k_i$ and the unique dispersion of polaritons allows for this process to take place in a resonant way, $2\varepsilon_{LPB}(k_p) \simeq \varepsilon_{LPB}(k_s) + \varepsilon_{LPB}(k_i)$. The onset of parametric oscillation is clearly visible in the emission pattern for pump intensities above the threshold $I_p^{th} = 12.5$ mW (upper panel of Fig.1): the occupation of the signal at $k_s \simeq 0$ becomes in this case very large and the linewidth of the emission in energy is substantially reduced as compared to the bare LPB linewidth shown in the lower panel. The spectral narrowing in energy is accompanied by a significant broadening of the k-space emission. The flat shape of the coherent OPO emission in the $(k_x; E)$ plane is however more likely to be a consequence of the peculiar shape of finite-size non-equilibrium condensates discussed in **(author?)** [15] rather than an evidence of the diffusive nature of the Goldstone mode **(author?)** [16].



Figure 1: Direct experimental observation on the CCD camera of a two-dimensional section $(k_x; E)$, with $k_y = 0$, of the lower polariton dispersion for pump powers I_p (a) just above (13 mW) and (b) below (12 mW) the OPO threshold $I_p^{th} = 12.5$ mW. Energy spectra at $k_x = 0$ (dashed line in the figure) are depicted on the right, with a magnification of a factor 3 for the lower panel of the figure. The PL emission is normalised to 1 and plotted in a linear color scale. (color online)

To clearly identify the threshold, we have studied the energy of the signal emission as a function of pump power (full triangles in Fig.2). A smooth and almost linear blue-shift of the signal energy is visible at low pump powers, while a sudden jump appears for I_p just above 12 mW due to the onset of parametric oscillation. Such a discon-



Figure 2: Energy (full triangles) and decay time (open circles) of the signal versus pump power. The red dashed line indicates the pump threshold. Inset: time evolution of the energy-integrated signal emission ΔI_s for three pump powers (1 mW, 7.5 mW, 12 mW). (color online)

tinuous behaviour around the threshold was predicted in Ref **(author?)** [14]**(author?)** [10]: below the threshold, the energy of the (incoherent) parametric luminescence is indeed fixed by the slightly blue-shifted LPB dispersion, while above threshold it is determined by a more complex OPO dynamics that also involves the idler energy. Far above threshold, the blue-shift saturates.

The main object of the present study is the response of the system in its stationary state to an additional weak ($\lesssim 0.2\,\mathrm{mW}$) probe pulse that impinges on the sample at a large angle of around 20°, i.e. in the vicinity of the idler. The evolution of the system in response to the probe pulse is monitored by investigating the time- and momentum-resolved signal emission and, in particular, its decay time. As an example, we have traced in the inset of Fig.2 the time-evolution of the difference $\Delta I_s = I_s(pump + probe) - I_s(pump)$ between the signal emission intensity in the presence and in the absence of the probe, respectively, for three different pump intensity values. To rule out non-linear effects and ensure we are in a linear-response regime with respect to the probe intensity, we have checked that the physics of interest is independent of the probe pulse intensity: while some changes remain visible in the short-time dynamics, the long-time dynamics simply shows a global rescaling of the observed intensity.

Right after the arrival of the probe pulse, parametric scattering of pump polaritons into the k_s state is stimulated by the small population of the new idler polaritons injected by the probe: in the plotted curves, this corresponds to a fast switch-on of the ΔI_s at the probe arrival time. The fast decay on a 30 ps scale is then followed by a much slower exponential decay on a time scale in the 100 ps range, i.e. orders of magnitude longer than both the empty cavity decay time (2 ps) and the polariton-polariton scattering time **(author?)** [6], but still significantly shorter than the probe repetition time of approximately 12 ns).



Figure 3: k_x dependence of the signal decay time for two different pump powers, 11.5 mW (open circles) and 12.5 mW (full circles). The high values of the decay time ($> 500\,\mathrm{ps}$) far from $k_x = 0$ are due to the integration over all the spot area and over all the emitted energies: the intensity profile (not shown) for both 11.5 mW and 12.5 mW is strongly peaked in the parametric scattering region around the signal state at $k_x \simeq 0$, while a much weaker and slower incoherent emission coming from higher energy states (e.g. residual excitons) and from the border of the spot is responsible for the tails at $|k_x| > 0.5\,\mu\mathrm{m}^{-1}$. (color online)

As easily seen by comparing the three curves shown in the inset, the response of the system strongly depends on the intensity I_p of the cw pump. While the decay time of the transient decreases for increasing pump powers and eventually goes below the streak-camera resolution of $\backsim 30$ ps, the long decay-time significantly increases with I_p. This latter dependence is summarized by the open circles in the main panel of Fig.2. The decay time shows a divergent behavior **(author?)** [17] for pump powers approaching $I_p = 12.5\,\mathrm{mW}$; for higher pump powers, it exceeds the time window of our setup. The parametric nature of the enhanced life time is confirmed by the coincidence of the divergence with the signal emission energy jump and the frequency narrowing of the luminescence (see Fig. 1).

The role of the parametric processes is further evidenced by the momentum-resolved data shown in Fig.3, in which the decay time of the different k_s components of the signal emission is plotted as a function of the wavevector k_x. The considered wavevector range is centered around the value k_s where signal emission would appear if the pump intensity was above threshold. While the decay time is a smooth function of k_x for I_p well below the parametric oscillation threshold (open circles), a marked peak is apparent in the vicinity of k_s ($k_x \approx 0$) for pump intensities around and above the threshold (full circles).

A convenient way to interpret the observed slowing-down of the response to the probe is to use the coherent polariton model based on a pair of coupled Gross-Pitaevskii-like nonlinear wave equations for respectively the photon and exciton fields $\psi_{C,X}(\mathbf{r},t)$ **(author?)** [18, 19],

$$i\,\frac{\partial \psi_C}{\partial t} = \left(\omega_C(-i\nabla_{\mathbf{r}}) - i\frac{\gamma_C}{2}\right)\psi_C + \Omega_R\,\psi_X + F(\mathbf{r},t) \quad (1)$$

$$i\,\frac{\partial \psi_X}{\partial t} = \left(\omega_X\,\psi_X - i\frac{\gamma_X}{2}\right)\psi_X + \Omega_R\,\psi_C + g\,|\psi_X|^2\,\psi_X \quad (2)$$

We follow the dynamics of the system starting from the $\psi_{X,C}(\mathbf{r},t) = 0$ vacuum state. $\omega_C(\mathbf{k})$ is the photon dispersion, while the exciton dispersion is assumed to be flat at ω_X. $\gamma_{C,X}$ are the decay rates of the cavity-photon and the exciton, respectively. The exciton-exciton interactions are characterized by the nonlinear coupling coefficient g and Ω_R is the exciton-photon Rabi coupling. The driving $F(\mathbf{r},t)$ is proportional to the incident electromagnetic field and has to include both the CW pump and the pulse probe: once the system has attained its stationary-state under the CW-pump only, an additional short probe pulse is applied close to resonance with the idler. The following response of the system is monitored on the most relevant observables, in particular the polariton distribution in k-space. For the sake of simplicity, we have limited ourselves to the case of a plane-wave pump with a well-defined wavevector k_p and periodic boundary conditions, while the finite spatial size of the probe beam is fully taken into account.

As discussed in Ref. **(author?)** [10, 19], the approach to the OPO threshold from below is signalled by the decay rate of some mode tending to zero. As a function of k, the decay time results then strongly peaked around the point where the decay rate is the smallest: the closer to threshold, the higher the peak value. These general claims are perfectly visible in the numerical result plotted in Fig.4. For pump intensities just below I_p^{th}, the evolution of the ntegrated signal emission $\Delta I_s(t)$ after the arrival of the probe pulse is characterized by a short transient followed by a much slower exponential decay, with a time constant that dramatically increases as the threshold is approached [panel (a)]. By comparing the overall decay time of the integrated $\Delta I_s(t)$ [panel (b)] with the k-dependent decay time [panel (c)], it is immediate to see that the former is determined by the decay time of the longest-lived mode, a quantity that increases in magnitude and becomes progressively more peaked as the threshold is approached.



Figure 4: Results of numerical calculations. Upper (a) panel: time dependence of k-integrated signal emission ΔI_s for different values of I_p; integration is performed over the k-space region surrounding k_s. Central (b) panel: decay time of ΔI_s as a function of CW pump intensity. Lower (c) panel: k-dependence of decay rate for different values of $I_p/I_p^{th} = 0.57$, 0.95, 0.99 below threshold. Inset (d): k-dependence of the decay time for $I_p/I_p^{th} = 1.15$, above threshold; the vertical dotted line indicates the coherent signal emission wavevector k_s, at which the decay time is infinite by definition of spontaneous symmetry breaking. (color online)

This theoretical result is in good agreement with the experimental observations for pump intensities in the vicinity of the threshold, but some specific attention has to be paid at the experimental data for very low pump power. In this regime, the theoretical calculations predict that the decay time should go back to the bare polariton lifetime, while a quite long decay time is observed in the experiment even well below the oscillation threshold and at wavevectors far from the signal emission. To explain this behaviour, one can mention the spatial inhomogeneity of the system that smoothens the k-space distribution, as well as the presence of residual excitons that have accumulated into long-lived states and that relax down on a long time scale. Clearly, these incoherent scattering processes are most important for low pump powers, while coherent parametric processes take it over as the threshold is approached.

As we already mentioned, the decay time above threshold is too long to be quantitative measured with the present setup. Numerical simulations do not suffer from such a difficulty, and we summarize here the main features that one expects for this regime. As a consequence of the spontaneously broken $U(1)$ symmetry, the spectrum of the elementary excitations is characterized by the presence of a soft Goldstone mode: as the wavevector $q = k - k_s$ of the excitation tends to zero, both its frequency and decay rate tend to zero **(author?)** [16]. This prediction is confirmed by the numerical results for the k-dependent decay rate that we show in Fig.4(d): once again, the peaked structure of the decay rate as a function of $k - k_s$ is apparent. It should be mentioned that, even though the long decay time of the slowest decaying mode eventually appears in the very long time behaviour of the integrated $\Delta I_s(t)$, this feature is often masked by other, faster decaying branches that contribute in a much more substantial way to the observed ΔI_s signal.

In conclusion, we have investigated the response of a continuously pumped optical parametric oscillator to an additional weak pulse in the vicinity of the idler. The emission from the signal is resolved in time and its momentum distribution is analyzed as a function of the pump intensity: when the pump intensity is close or above the parametric oscillation threshold, a lifetime orders of magnitude longer than the intrinsic polariton lifetime is observed for polariton modes in the vicinity of the signal emission. The experimental observations are explained in terms of a critical slowing down of the elementary excitation dynamics as the threshold is approached, and then of the presence of a soft Goldstone mode above the threshold.

This work was partially supported by the Spanish MEC (MAT2005-01388, NAN2004-09109-C04-04 & QOIT-CSD2006-00019) and the CAM (S-0505/ESP-0200). D.B. acknowledges a scholarship of the FPU program of the Spanish MEC and D.S. thanks the Ramon y Cajal Program. I.C. is indebted to C. Ciuti for continuous discussions and acknowledges support from the italian MIUR, the french CNRS and IFRAF and the EuroQUAM-FerMix program.

---

[1] D. Pines and P. Nozieres, *The Theory of Quantum Liquids*, Vols. 1 and 2 (Addison-Wesley, Redwood City, 1966); L. Pitaevskii and S. Stringari, *Bose Einstein Condensation* (Oxford University Press, Oxford, 2003).

[2] *Phase transitions and critical phenomena*, vol.17 (Statistical mechanics of driven diffusive systems), eds. C.Domb and J.L.Lebowitz, Academic, NY, 1989

[3] Semicond. Sci. Technol. **18**, *Special Issue on Microcavities*, Guest Editors J. Baumberg and L. Viña, Publisher S. Quin (Bristol, UK, 2003).

[4] J. Keeling, F. M. Marchetti, M. H. Szymanska, P. B. Littlewood, Semicond. Sci. Technol. **22** R1 (2007).

[5] R. M. Stevenson, *et al.*, Phys. Rev. Lett. **85**, 3680 (2000); J. J. Baumberg, *et al.*, Phys. Rev. B **62**, R16247 (2000); R. Houdré *et al.*, Phys. Rev. Lett. **85**, 2793 (2000); A. Baas *et al.* Phys. Rev. Lett. **96**, 176401 (2006).

[6] A. Huynh, *et al.*, Phys. Rev. Lett. **90**, 106401 (2003).

[7] N. A. Gippius *et al.*, Europhys. Lett. **67**, 997 (2004); D. N. Krizhanovskii *et al.*, Phys. Rev. B **77**, 115336 (2008).

[8] D. D. Solnyshkov, I. A. Shelykh, N. A. Gippius, A. V. Kavokin, G. Malpuech, Phys. Rev. B 77, 045314 (2008)

[9] J. Kasprzak *et al.*, Nature **443**, 409 (2006); H. Deng *et al.*, Phys. Rev. Lett. **99**, 126403 (2007); R. Balili *et al.*, Science **316**, 1007 (2007).

[10] M.Wouters, I.Carusotto, Phys. Rev. B **75**, 075332 (2007).

[11] I.Carusotto and C.Ciuti, Phys.Rev.B **72**, 125335 (2005).

[12] K. Huang, *Statistical Mechanics* (Wiley, NY, 1963).

[13] M. C. Cross and P. C. Hohenberg, Rev. Mod. Phys. **65**, 851 (1993).

[14] D. M. Whittaker, Phys. Rev. B **71**, 115301 (2005).

[15] M. Wouters, I. Carusotto, and C. Ciuti, Phys. Rev. B **77**, 115340 (2008).

[16] M.Wouters, I.Carusotto, Phys.Rev. A **76**, 043807 (2007).

[17] Such long-time features were not accessible to frequency-domain luminescence experiments, see e.g. P.G. Savvidis, *et al.*, Phys. Rev. B **64**, 075311 (2001).

[18] C. Ciuti, P. Schwendimann, and A. Quattropani, Semicond. Sci. Technol. **18**, S279 S293 (2003).

[19] C. Ciuti and I. Carusotto, Phys. Stat. Sol. (b) **242**, 2224 (2005).